



Acknowledgments:

We would like to express our gratitude to the authors whose works have been arranged in this booklet: their insights and expertise greatly assisted this prime selection.

We are dedicated to developing and providing high-quality and cutting-edge biological sample collection products for human genomics, infectious diseases, environmental and forensic applications, along with automated workflow solutions. Copan's innovative approach enables an ever-expanding community of laboratories, scientists, and institutions to benefit from an accessible sample collection that guarantees reliable quality performance.

Our goal is to continue this innovation by providing products, customized services, and prime solutions to improve patients' health and wellness.

In this booklet, you'll find a selection of the most interesting and recent independent studies where our products designed specifically for self-collection are used. By reading them, you'll discover how self-collection offers many advantages, such as lowering cultural barriers that limit testing and increasing participation in clinical studies and screening programs while simultaneously saving time and resources in hospitals and POC.

A patented technology

Why so special?

Copan conceived the FLOQ® technology in 2003, to answer the need of healthcare professionals for a more efficient sample collection. The main feature of this technology is the perpendicular arrangement of the short Nylon® fibers. This invention revolutionized the preanalytical world, bringing sampling to the next level.



FLOQSwabs®

FLOQSwabs® are made of a solid molded plastic applicator shaft with a tip, which they can both vary in size and shape. Thanks to their patented flocked tip, FLOQSwabs® ensure a flawless specimen collection, which expands downstream diagnostic testing capabilities.

Media pairing

The perfect match

The combination of FLOQSwabs® with different Bacteriology and Virology transport media expands their flexibility even further.

Whether you have a specific collection site, pathogen, or downstream application in mind, Copan will be able to suggest the best combination to suit your needs.

Look for the mark in all our printed material and online content!





Self Vaginal FLOQSwabs®

The FLOQSwabs® version we designed for vaginal home-collection. With their soft and small tip, easy handling, insertion length mark, and clear instructions, Self Vaginal FLOQSwabs® will make every woman forget unpleasant tests. Also, being dry translates into cost-effective transport without any liquid-related safety issues!



Self UriSponge™

Self UriSponge™ is our safe and easy-to-use device to collect urine samples at home. The sponge absorbs the correct amount of first-void urine sample without risk and discomfort, while the leak-proof tube ensures safe handling and shipment. Thanks to the preservatives and its formula, Self UriSponge™ allows the shipment of urine samples from remote areas and fits with the main molecular platforms.

Self-Collection

An Observational Study Comparing HPV Prevalence and Type Distribution Between HPV-Vaccinated and -Unvaccinated Girls After Introduction of School-Based HPV Vaccination in Norway

Espen Enerly et al.

PLoS One. 2019 Oct 10;14(10):e0223612

9

STOP HPV Study Protocol: a Nationwide Case–Control Study of the Association Between Oropharyngeal Cancer and Human Papillomavirus (HPV) Infection in Brazil

Eliana Marcia Wendland et al.

BMJ Open. 2020 Jan 29;10(1):e031602

10

Analytical Performance of HPV Assays on Vaginal Self-Collected vs Practitioner-Collected Cervical Samples: the SCoPE Study

M. Saville et al.

J Clin Virol. 2020 Jun;127:104375

11

Population Impact of Girls-Only Human Papillomavirus 16/18 Vaccination in The Netherlands: Cross-Protective and Second-Order Herd Effects

Joske Hoes et al.

Clin Infect Dis. 2021 Mar 1;72(5):e103-e111

12

Treating Male Partners Of Women with Bacterial Vaginosis (Stepup): a Protocol for a Randomised Controlled Trial to Assess the Clinical Effectiveness of Male Partner Treatment for Reducing the Risk of Bv Recurrence

Lenka A. Vodstrcil et al.

BMC Infect Dis. 2020 Nov 11;20(1):834

13

Assessing the Impact of Mailing Self-Sampling Kits for Human Papillomavirus Testing to Unscreened Non-Responder Women in Manitoba

F. Jalili et al.

Curr Oncol. 2019 Jun;26(3):167-172

14

Self-Collection for Under-Screened Women in a National Cervical Screening Program: Pilot Study

M. Saville et al.

Curr Oncol. 2018 Feb;25(1):e27-e32

15

A comparison of cotton and flocked swabs for vaginal self-sample collection

M. Viviano et al.

Int J Womens Health. 2018 May 15;10:229-236

16

Self-Collection

Randomized Comparison of Two Vaginal Self-Sampling Methods for Human Papillomavirus Detection: Dry Swab versus FTA Cartridge

R. Catarino et al.

PLoS One. 2015 Dec 2;10(12):e0143644

17

The acceptability and cost of a home-based chlamydia retesting strategy: findings from the REACT randomised controlled trial

K.S. Smith et al.

BMC Public Health. 2016 Jan 28;16:83

18

UriSwab: an effective transport medium for nucleic acid detection of *Chlamydia trachomatis*, *Mycoplasma genitalium* and *Neisseria gonorrhoeae*

A. M. G. Costa et al.

Sex Health. 2017 Nov;14(6):502-506

19

A Randomized Comparison of Different Vaginal Self-sampling Devices and Urine for Human Papillomavirus Testing—Predictors 5.1

Louise Cadman et al.

Cancer Epidemiol Biomarkers Prev. 2021 Apr;30(4):661-668.

20

The feasibility and acceptability of self-sampling and HPV testing using Cepheid Xpert® HPV in a busy primary care facility

YL Woo

J Virus Erad. 2019 Mar 4;5(Suppl 1):10-11.

21

Randomized trial evaluating self-sampling for HPV DNA based tests for cervical cancer screening in Nigeria

Fatima Modibbo et al.

Infect Agent Cancer. 2017; 12: 11.

22

Acceptability of human papillomavirus (HPV) self-sampling among never- and under-screened Indigenous and other minority women: a randomised three-arm community trial in Aotearoa New Zealand

Naomi Brewer et al.

Lancet Reg Health West Pac. 2021 Sep 7;16:100265

23

The Acceptability and Preference of Vaginal Self-sampling for Human Papillomavirus (HPV) Testing among a Multi-ethnic Asian Female Population

Su Pei Khoo et al.

Cancer Prev Res (Phila). 2021 Jan;14(1):105-112.

24

Self-Collection

Implementing the 3T-approach for cervical cancer screening in Cameroon: Preliminary results on program performance

Juliette Levy1 et al.

Cancer Med. 2020 Oct;9(19):7293-7300.

25

Exploring Factors Associated with Patients Who Prefer Clinician-Sampling to HPV Self-Sampling: A Study Conducted in a Low-Resource Setting

Jessica Sormani et al.

Int J Environ Res Public Health. 2022 Jan; 19(1): 54.

26

Non-speculum sampling approaches for cervical screening in older women: randomised controlled trial

Rebecca Landy et al.

Br J Gen Pract. 2021 Dec 31;72(714):e26-e33.

27

Comparison of Different Self Sampling Devices for Molecular Detection of Human Papillomavirus (HPV) and Other Sexually Transmitted Infections (STIs): A Pilot Study

Illari Sechi et al.

Healthcare (Basel). 2022 Feb 28;10(3):459.

28

Saliva detection of SARS-CoV-2 for mitigating company outbreaks: a surveillance experience, Milan, Italy, March 2021

Emerenziana Ottaviano et al.

Epidemiol Infect. 2021 Jul 30;149:e171.

29

Self-Collection

An Observational Study Comparing HPV Prevalence and Type Distribution Between HPV-Vaccinated and -Unvaccinated Girls After Introduction of School-Based HPV Vaccination in Norway



Espen Enerly¹, Ragnhild Flingsborg¹, Irene Kraus Christiansen², Suzanne Campbell¹, Mona Hansen², Tor Åge Myklebust^{3,4}, Elisabeth Weiderpass⁵, Mari Nygård¹

Affiliation: ¹Department of Research, Cancer Registry of Norway, Oslo, Norway ²Department of Microbiology and Infection Control, National HPV Reference Laboratory, Akershus University Hospital, Lørenskog, Norway ³Department of Registration, Cancer Registry of Norway, Oslo, Norway ⁴Department of Research and Innovation, Møre and Romsdal Hospital Trust, Ålesund, Norway ⁵International Agency for Research on Cancer (IARC), World Health Organization, Lyon, France

Keywords

FLOQSwabs®

Oral Sampling

HPV

HPV Vaccination

Abstract

Background: Many countries have initiated school-based human papillomavirus (HPV) vaccination programs. The real-life effectiveness of HPV vaccines has become increasingly evident, especially among girls vaccinated before HPV exposure in countries with high vaccine uptake. In 2009, Norway initiated a school-based HPV vaccination program for 12-year-old girls using the quadrivalent HPV vaccine (Gardasil®), which targets HPV6, 11, 16, and 18. Here, we aim to assess type-specific vaginal and oral HPV prevalence in vaccinated compared with unvaccinated girls in the first birth cohort eligible for school-based vaccination (born in 1997).

Methods: This observational, cross-sectional study measured the HPV prevalence ratio (PR) between vaccinated and unvaccinated girls in Norway. Facebook advertisement was used to recruit participants and disseminate information about the study. Participants self-sampled vaginal and oral specimens using an Evalyn® Brush and a FLOQSwab®, respectively. Sexual behaviour was ascertained through a short questionnaire.

Results: Among the 312 participants, 239 (76.6%) had received at least one dose of HPV vaccine prior to sexual debut. 39.1% of vaginal samples were positive for any HPV type, with similar prevalence among vaccinated and unvaccinated girls (38.5% vs 41.1%, PR: 0.93, 95% confidence interval [CI]: 0.62–1.41). For vaccine-targeted types there was some evidence of lower prevalence in the vaccinated (0.4%) compared to the unvaccinated (6.8%) group (PR: 0.06, 95%CI: 0.01–0.52). This difference remained after adjusting for sexual behaviour (PR: 0.04, 95%CI: 0.00–0.42). Only four oral samples were positive for any HPV type, and all of these participants had received at least one dose of HPV vaccine at least 1 year before oral sexual debut.

Conclusion: There is evidence of a lower prevalence of vaccine-targeted HPV types in the vagina of vaccinated girls from the first birth cohort eligible for school-based HPV vaccination in Norway; this was not the case when considering all HPV types or types not included in the quadrivalent HPV vaccine.

Self-Collection

STOP HPV Study Protocol: a Nationwide Case–Control Study of the Association Between Oropharyngeal Cancer and Human Papillomavirus (HPV) Infection in Brazil



Eliana Marcia Wendland^{1,2}, Natalia Luiza Kops¹, Juliana Comerlato¹, Jaqueline Driemeyer Correia Horvath¹, Marina Bessel¹, Daniel Sperb³, Cristina Pimenta⁴, Flávia Moreno Alves de Souza⁴, Gerson Fernando Mendes Pereira⁴, Frederico Soares Falcetta¹

Affiliation: ¹Escritório de Projetos PROADI-SUS, Hospital Moinhos de Vento, Porto Alegre, Rio Grande do Sul, Brazil

²Universidade Federal de Ciências da Saúde de Porto Alegre, Porto Alegre, Rio Grande do Sul, Brazil ³Hospital Moinhos de Vento, Porto Alegre, Rio Grande do Sul, Brazil ⁴Department of STIs, AIDS and Viral Hepatitis, Ministry of Health, Brasília, Brazil

Keywords

FLOQSwabs®

HPV

Genital Sampling

Abstract

Introduction: Human papillomavirus (HPV) is the most common sexually transmitted infection and is associated with several types of cancer. The number of cases of HPV-associated head and neck squamous cell carcinomas (HNSCCs), especially oropharyngeal carcinomas, has increased significantly in recent years despite decreased tobacco smoking rates. Currently, no data concerning the risk factors and prevalence of HPV in HNSCC patients in all regions of Brazil are available, making it difficult to promote advances in this field of public health. Therefore, our goal is to determine the impact of infection by HPV, including HPVs with different genotypes, on head and neck cancer and the risk factors associated with the development of head and neck cancer in all regions of Brazil.

Methods and analysis: This is a case–control study that will include 622 patients and 622 controls from all regions of Brazil. A questionnaire will be applied to gather information on sociodemographic, behavioural and health factors. Oral, cervical or penile/scrotal, and anal specimens and serum samples will be collected from all participants with Copan FLOQSwabs®. Formalin-fixed paraffin-embedded tissue from tumour biopsies will be analysed only in the case group. Molecular and serological analyses will be performed to evaluate the presence and role of HPV in the development of head and neck cancer.

Results: The results will provide a better understanding of the association between HPV infection and head and neck cancer, allowing the identification of at-risk individuals and generating hypotheses for the use of preventive or screening measures in specific population groups.

Self-Collection

Analytical Performance of HPV Assays on Vaginal Self-Collected vs Practitioner-Collected Cervical Samples: the SCoPE Study



M. Saville^{1,2,3,4}, D. Hawkes^{1,2,5,6}, MHT Keung^{1,2}, ELO Ip^{1,2}, J. Silvers⁷, F. Sultana^{1,8}, MJ Malloy^{1,8}, LS Velentzis^{8,9}, K. Canfel L.^{9,10}, CD Wrede^{3,7}, JML Brotherton^{1,4,8}

Affiliation: ¹VCS Foundation, Carlton South, Victoria, Australia ²VCS Pathology, Carlton South, Victoria, Australia ³Department of Obstetrics and Gynaecology, University of Melbourne, Parkville, Victoria, Australia ⁴Department of Obstetrics and Gynaecology, University of Malaya, Kuala Lumpur, Malaysia ⁵Department of Pharmacology and Therapeutics, University of Melbourne, Parkville, Victoria, Australia ⁶Department of Pathology, University of Malaya, Kuala Lumpur, Malaysia ⁷Department of Oncology and Dysplasia, Royal Women's Hospital, Melbourne, Victoria, Australia ⁸Melbourne School of Population and Global Health, University of Melbourne, Parkville, Victoria, Australia ⁹Cancer Research Division, Cancer Council NSW, Woolloomooloo, NSW, Australia ¹⁰School of Public Health, Sydney Medical School, University of Sydney, NSW, Australia

Keywords

FLOQSwabs®

Vaginal Sampling

HPV

Diagnostics Platforms

Abstract

Background: In the last decade, human papillomavirus (HPV) testing has been evaluated extensively for cervical screening, with studies finding increased sensitivity compared to cytology. Another advantage of HPV based-screening is the ability to test vaginal samples that can be collected by women themselves. Self-collection has the potential to extend cervical screening coverage by increasing participation rates, particularly among women who are under-screened or have never screened. This could have a significant impact on cervical cancer prevention, as the majority of invasive cervical cancer cases occur among under-screened women. Both the Netherlands and Australia have transitioned their national programs from cytology to HPV as the primary screening test and both countries include a pathway for self-collection.

Objectives: We evaluated the relative sensitivity for HPV detection of self-collection compared with practitioner-collected cervical specimens in the context of the Australian National Cervical Screening Program (NCSP). **Study Design:** 303 women aged ≥ 18 years attending a single tertiary referral centre took their own sample using a Copan FLOQSwabs®, and then had a practitioner-collected sample taken at colposcopy. All samples were tested at a single laboratory on the six PCR-based HPV assays which can be utilised in the NCSP; Roche cobas 4800 and cobas, Abbott Real Time, BD Onclarity, Cepheid Xpert, and Seegene Anyplex.

Results: HPV16/18 results had high observed agreement between self- and practitioner-collected samples on all assays (range: 0.94-0.99), with good agreement for non-HPV16/18 oncogenic HPV types (range: 0.64-0.73).

Conclusions: Self-collection for HPV-based cervical screening shows good concordance and relative sensitivity when compared to practitioner collected samples across assays in the NCSP.

Self-Collection

Population Impact of Girls-Only Human Papillomavirus 16/18 Vaccination in The Netherlands: Cross-Protective and Second-Order Herd Effects



Joske Hoes^{1,2}, Petra J. Woestenberg^{1,3}, Johannes A. Bogaards^{1,2}, Audrey J. King¹, Hester E. de Melker¹, Johannes Berkhof², Christian J. P. A. Hoebe^{3,4}, Marianne A. B. van der Sande^{5,6}, Birgit H. B. van Benthem¹

Affiliation: ¹Centre for Infectious Disease Control, National Institute for Public Health and the Environment, Bilthoven, The Netherlands ²Department of Epidemiology and Data Science, Amsterdam University Medical Center, location VUmc, Amsterdam, The Netherlands ³Department of Social Medicine, Care and Public Health Research Institute, Maastricht University Medical Center, Maastricht, The Netherlands ⁴Department of Sexual Health, Infectious Diseases and Environment, South Limburg Public Health Service, Heerlen, The Netherlands ⁵Department of Public Health, Institute of Tropical Medicine, Antwerp, Belgium ⁶Julius Center, University Medical Center Utrecht, Utrecht, The Netherlands

Keywords

FLOQSwabs®

HPV

Vaccination

Genital Sampling

Abstract

Background: Human papillomavirus (HPV) vaccination programs achieve substantial population-level impact, with effects extending beyond protection of vaccinated individuals. We assessed trends in HPV prevalence up to 8 years postvaccination among men and women in the Netherlands, where bivalent HPV vaccination, targeting HPV types 16/18, has been offered to (pre)adolescent girls since 2009 with moderate vaccination coverage.

Methods. We used data from the PASSYON study, a survey initiated in 2009 (pre-vaccination) and repeated biennially among 16- to 24 year-old visitors of sexual health centers obtained starting from genital self-collection swab (Copan FLOQSwabs®). We studied genital HPV positivity from 2009 to 2017 among women, heterosexual men, and unvaccinated women using Poisson generalized estimating equation models, adjusted for individual- and population-level confounders. Trends were studied for 25 HPV types detected by the SPF10-LiPA25 platform.

Results. A total of 6354 women (64.7% self-reported unvaccinated) and 2414 heterosexual men were included. Percentual declines in vaccine types HPV-16/18 were observed for all women (12.6% per year [95% confidence interval {CI}, 10.6–14.5]), heterosexual men (13.0% per year [95% CI, 8.3–17.5]), and unvaccinated women (5.4% per year [95% CI, 2.9–7.8]). We observed significant declines in HPV-31 (all women and heterosexual men), HPV-45 (all women), and in all high-risk HPV types pooled (all women and heterosexual men). Significant increases were observed for HPV-56 (all women) and HPV-52 (unvaccinated women).

Conclusions. Our results provide evidence for first-order herd effects among heterosexual men against HPV-16/18 and cross protective types. Additionally, we show second-order herd effects against vaccine types among unvaccinated women. These results are promising regarding population-level and clinical impact of girls-only bivalent HPV vaccination in a country with moderate vaccine uptake.

Self-Collection

Treating Male Partners of Women with Bacterial Vaginosis (Stepup): a Protocol for a Randomised Controlled Trial to Assess the Clinical Effectiveness of Male Partner Treatment for Reducing the Risk of Bv Recurrence



Lenka A. Vodstrcil^{1,2}, Erica L. Plummer^{1,2}, Michelle Doyle², Christopher K. Fairley^{1,2}, Colette McGuinness², Deborah Bateson^{3,4}, Jane S. Hocking⁵, Matthew G. Law⁶, Kathy Petoumenos⁶, Basil Donovan⁶, Eric P. F. Chow^{1,2}, Catriona S. Bradshaw^{1,2,5} and on behalf of the StepUp RCT Team

Affiliation: ¹Central Clinical School, Monash University, Carlton, VIC 3053, Australia ²Melbourne Sexual Health Centre, Alfred Health, Carlton, VIC 3053, Australia ³Family Planning New South Wales, Ashfield 2131, Australia ⁴Discipline of Obstetrics, Gynaecology and Neonatology, University of Sydney, Camperdown 2006, Australia ⁵Centre for Epidemiology and Biostatistics, Melbourne School of Population & Global Health, University of Melbourne, Parkville 3010, Australia ⁶The Kirby Institute, University of New South Wales, Sydney 2052, Australia

Keywords

FLOQSwabs®

Bacterial Vaginosis

Male Partners

Antibiotic Treatment

Abstract

Background: Bacterial vaginosis (BV) is estimated to affect 1 in 3 women globally and is associated with obstetric and gynaecological sequelae. Current recommended therapies have good short-term efficacy but 1 in 2 women experience BV recurrence within 6 months of treatment. Evidence of male carriage of BV-organisms suggests that male partners may be re-infecting women with BV-associated bacteria (henceforth referred to as BV-organisms) and impacting on the efficacy of treatment approaches solely directed to women. This trial aims to determine the effect of concurrent male partner treatment for preventing BV recurrence compared to current standard of care.

Methods: StepUp is an open-label, multicentre, parallel group randomised controlled trial based on genital self-collection (Copan FLOQSwabs®) for women diagnosed with BV and their male partner. Women with clinical-BV defined using current gold standard diagnosis methods (≥ 3 Amsel criteria and Nugent score (NS) = 4–10) and with a regular male partner will be assessed for eligibility, and couples will then be consented. All women will be prescribed oral metronidazole 400 mg twice daily (BID) for 7 days, or if contraindicated, a 7-day regimen of topical vaginal 2% clindamycin. Couples will be randomised 1:1 to either current standard of care (female treatment only), or female treatment and concurrent male partner treatment (7 days of combined antibiotics - oral metronidazole tablets 400 mg BID and 2% clindamycin cream applied topically to the glans penis and upper shaft [under the foreskin if uncircumcised] BID). Couples will be followed for up to 12 weeks to assess BV status in women, and assess the adherence, tolerability and acceptability of male partner treatment. The primary outcome is BV recurrence defined as ≥ 3 Amsel criteria and NS = 4–10 within 12 weeks of enrolment. The estimated sample size is 342 couples, to detect a 40% reduction in BV recurrence rates from 40% in the control group to 24% in the intervention group within 12 weeks.

Discussion: Current treatments directed solely to women result in unacceptably high rates of BV recurrence. If proven to be effective the findings from this trial will directly inform the development of new treatment strategies to impact on BV recurrence.

Self-Collection

Assessing the Impact of Mailing Self-Sampling Kits for Human Papillomavirus Testing to Unscreened Non-Responder Women in Manitoba



F. Jalili¹, C. O'Conaill¹, K. Templeton¹, R. Lotocki¹, G. Fischer², L. Manning², K. Cormier², K. Decker^{3,4}

Affiliation: ¹CervixCheck, CancerCare Manitoba, MB, Canada ²Diagnostic Services, Shared Health Manitoba, MB, Canada ³Department of Community Health Sciences, University of Manitoba, MB, Canada ⁴Research Institute in Oncology and Hematology, CancerCare Manitoba, Winnipeg, MB, Canada

Keywords

FLOQSwabs®

HPV

Cancer Screening

Mailing Self Sampling Kits

Abstract

Background: CervixCheck, Manitoba's cervical cancer screening program, conducted a pilot study to assess whether screening participation could be improved in unscreened women by offering a mailed self-sampling kit for human papillomavirus (HPV) testing instead of a Pap test.

Methods: In a prospective cohort study design, a sample of unscreened women (n = 1052) who had been sent an invitation letter from CervixCheck in the past but who did not respond were randomized to either an intervention group or a control group. The intervention group received a mailed HPV self-sampling kit with Copan FLOQSwabs®; the control group received no additional communication. Returned HPV self-sampling swabs were analyzed by a provincial laboratory. After 6 months, screening participation in the two study groups was compared using a logistic regression model adjusted for age and area of residence (urban or rural). Secondary outcomes included HPV positivity, specimen inadequacy, compliance with follow-up, and time to colposcopy.

Results: Screening participation was significantly higher in the intervention group than in the control group (n = 51, 9.6%, vs. n = 13, 2.5%; odds ratio: 4.7; 95% confidence interval: 2.56 to 8.77). Geographic area of residence (urban or rural) and age were not statistically significant.

Conclusions: The study demonstrated that HPV self-sampling kits can enhance screening participation in unscreened non-responder women in the setting of an organized screening program. Next steps should include additional research to determine the best implementation strategy for HPV self-sampling in Manitoba.

Self-Collection

Self-Collection for Under-Screened Women in a National Cervical Screening Program: Pilot Study



M. Saville¹, D Hawkes², E. Mclachlan³, S. Anderson⁴ K. Arabena³

Affiliation:¹Victorian Cytology Service, Carlton, Victoria, Australia; Department of Obstetrics and Gynaecology, University of Melbourne, Parkville, VIC 3053, Australia ²Victorian Cytology Service, Carlton, Victoria, Australia; Department of Pharmacology and Therapeutics, University of Melbourne, Parkville, VIC, Australia ³Indigenous Health Equity Unit, Melbourne School of Population and Global Health, University of Melbourne, Parkville, VIC, Australia ⁴Ballarat and District Aboriginal Collective, Baarlinjan Medical Clinic, Ballarat, VIC, Australia

Keywords

FLOQSwabs®

National Program

HPV

Cervical Cancer Screening

Abstract

Background: Commencing 1 December 2017, Australia introduced human papillomavirus (HPV)-based cervical screening. As part of this Australian renewed National Cervical Screening Program (NCSP) women who are either never- or under-screened and who refuse a practitioner collected sample will be able to collect their own sample for cervical screening. The aim of this study is to examine the quantitative results of a pilot study into the acceptability of the self-collection alternative pathway.

Methods: Eligible participants were offered the opportunity to collect their own sample. Those who agreed were given a flocked swab and an instruction sheet and took their own sample in an area of the health care clinic that afforded them adequate privacy. These samples were then given to clinic staff who returned them to Victorian Cytology Service (VCS) Pathology for HPV nucleic acid testing.

Results: Of 98 eligible women, seventy-nine undertook self-collection for HPV-based cervical screening. Seventyseven produced valid results, 14 were positive for oncogenic HPV, with 10 undertaking follow-up. Three women were found to have cervical squamous abnormalities with two of those being high-grade intraepithelial squamous lesions.

Conclusion: The pilot study for self-collection for cervical screening produced quantitative data that were similar to that already reported in the literature but had a much higher rate of acceptance compared with self-collection programs based in the home.

Self-Collection

A Comparison of Cotton and Flocked Swabs for Vaginal Self-Sample Collection



Manuela Viviano¹, Alexia Willame¹, Marie Cohen¹, Anne-Caroline Benski¹, Rosa Catarino¹, Christine Wuillemin¹, Phuong Lien Tran¹, Patrick Petignat¹, Pierre Vassilakos²
Affiliation: ¹Division of Gynecology, Geneva University Hospitals, Geneva, Switzerland. ²Geneva Foundation for Medical Education and Research, Geneva, Switzerland.

Keywords

FLOQSwabs®

HPV

Sample Release

Cellular Retrieval

Abstract

Objective: Vaginal self-sampling for human papillomavirus (HPV) testing has recently been proposed to optimize cervical cancer screening coverage. The objective of this study was to compare the performance of self-taken samples using flocked and cotton swabs for HPV detection and cellular retrieval.

Methods: We recruited women aged 21-65 years, referred to colposcopy at the Division of Gynecology of the Geneva University Hospitals between May and September 2016. Each participant collected 2 vaginal samples: 1 with a cotton swab (Copan ClassiqSwabs™) and 1 with a flocked swab (Copan FLOQSwabs®). A 1:1 randomization determined the order in which the 2 samples were taken. The swabs were introduced into a 20 mL PreservCyt® vial. Real-time polymerase chain reaction analysis using the Anyplex™ II HPV HR assay, cytofluorometric analysis and cytological cell counting were performed on each sample.

Results: A total of 119 participants were recruited in the study. Their mean \pm standard deviation age was 35.1 \pm 8.9 years. The HPV prevalence was 29.7% and 38.1% according to the cotton and flocked swab, respectively ($p=0.006$). The mean number of cells collected per milliliter according to cytofluorometry was 96,726.6 with the cotton swab and 425,544.3 with the flocked swab ($p<0.001$). The mean number of cells detected at cytological cell count was 13,130.42 using the cotton swab and 17,503.6 using the flocked swab ($p<0.001$).

Conclusion: The flocked swab achieved a greater cellular retrieval and showed an improved performance in HPV detection. Further studies are needed to assess the usability and cost-effectiveness of the 2 self-sampling devices.

Self-Collection

Randomized Comparison of Two Vaginal Self-Sampling Methods for Human Papillomavirus Detection: Dry Swab versus FTA Cartridge



Rosa Catarino^{1,2}, Pierre Vassilakos³, Aline Bilancioni¹, Mathieu Vanden Eynde¹, Ulrike Meyer-Hamme¹, Pierre-Alain Menoud⁴, Frédéric Guerry⁴, Patrick Petignat¹

Affiliation: ¹ Division of Gynecology, Department of Gynecology and Obstetrics, Geneva University Hospitals, Geneva, Switzerland. ² Faculty of Medicine, University of Geneva, Geneva, Switzerland. ³ Geneva Foundation for Medical Education and Research, Geneva, Switzerland. ⁴ Unilabs SA, Lausanne, Switzerland.

Keywords

FLOQSwabs®

Storage

HPV

Stability

Abstract

Background: Human papillomavirus (HPV) self-sampling (self-HPV) is valuable in cervical cancer screening. HPV testing is usually performed on physician-collected cervical smears stored in liquid-based medium. Dry filters and swabs are an alternative. We evaluated the adequacy of self-HPV using two dry storage and transport devices, the FTA cartridge and swab.

Methods: A total of 130 women performed two consecutive self-HPV samples. Randomization determined which of the two tests was performed first: self-HPV using Copan FLOQSwabs® (s-DRY) or vaginal specimen collection using a cytobrush applied to an FTA cartridge (s-FTA). After self-HPV, a physician collected a cervical sample using liquid-based medium (Dr-WET). HPV types were identified by real-time PCR. Agreement between collection methods was measured using the kappa statistic.

Results: HPV prevalence for high-risk types was 62.3% (95%CI: 53.7–70.2) detected by s-DRY, 56.2% (95%CI: 47.6–64.4) by Dr-WET, and 54.6% (95%CI: 46.1–62.9) by s-FTA. There was overall agreement of 70.8% between s-FTA and s-DRY samples (kappa = 0.34), and of 82.3% between self-HPV and Dr-WET samples (kappa = 0.56). Detection sensitivities for low-grade squamous intraepithelial lesion or worse (LSIL+) were: 64.0% (95%CI: 44.5–79.8) for s-FTA, 84.6% (95%CI: 66.5–93.9) for s-DRY, and 76.9% (95%CI: 58.0–89.0) for Dr-WET. The preferred self-collection method among patients was s-DRY (40.8% vs. 15.4%). Regarding costs, FTA card was five times more expensive than the swab (~5 US dollars (USD)/per card vs. ~1 USD/per swab).

Conclusion: Self-HPV using dry swabs is sensitive for detecting LSIL+ and less expensive than s-FTA.

Self-Collection

The Acceptability and Cost of a Home-Based Chlamydia Retesting Strategy: Findings from the REACT Randomised Controlled Trial



K. S. Smith¹, J. M. Kaldor¹, J. S. Hocking², M. S. Jamil¹, A. M. McNulty^{3,4}, P. Read⁵, C. S. Bradshaw^{6,7}, M. Y. Chen^{6,7}, C. K. Fairley^{6,7}, H. Wand¹, K. Worthington⁶, S. Blake³, V. Knight³, W. Rawlinson⁸, M. Saville⁹, S. N. Tabrizi^{10,11,12,13}, S. M. Garland^{10,11,12,13}, B. Donovan¹, R. Guy¹ Affiliation: ¹The Kirby Institute, UNSW Australia, Sydney, Australia ²Melbourne School of Population and Global Health, University of Melbourne, Melbourne, Australia ³Sydney Sexual Health Centre, Sydney, Australia ⁴School of Public Health and Community Medicine, UNSW Australia, Sydney, Australia ⁵Kirketon Road Centre, Sydney, Australia ⁶Melbourne Sexual Health Centre, Melbourne, Australia ⁷Central Clinical School, Monash University, Melbourne, Australia ⁸Serology and Virology Division, (SAViD) SEALS Microbiology, Prince of Wales Hospital, Sydney, Australia ⁹VCS Pathology, Melbourne, Australia ¹⁰Department of Obstetrics and Gynaecology, University of Melbourne, Melbourne, Australia ¹¹Department of Microbiology, Royal Children's Hospital, Melbourne, Australia. ¹²Department of Microbiology and Infectious Diseases, Royal Women's Hospital, Melbourne, Australia ¹³Murdoch Childrens Research Institute, Melbourne, Australia

Keywords

UriSwab™

Chlamydia trachomatis

Cost and Acceptability

Retesting

Abstract

Background: Chlamydia retesting three months after treatment is recommended to detect reinfections, but retesting rates are typically low. The REACT (retest after Chlamydia trachomatis) randomised trial demonstrated that home-based retesting using postal home-collection kits that included Copan UriSwab™ for urine self-collection and SMS reminders, resulted in substantial improvements in retesting rates in women, heterosexual men and men who have sex with men (MSM), with detection of more repeat positive tests compared with SMS reminder alone. In the context of this trial, the acceptability of the home-based strategy was evaluated and the costs of the two strategies were compared. **Methods:** REACT participants (200 women, 200 heterosexual men, 200 MSM) were asked to complete an online survey that included home-testing acceptability and preferred methods of retesting. The demographics, sexual behaviour and acceptability of home collection were compared between those preferring home-testing versus clinic-based retesting or no preference, using a chi-square test. The costs to the health system of the clinic-based and home retesting strategies and the cost per infection for each were also compared.

Results: Overall 445/600 (74 %) participants completed the survey; 236/445 from the home-testing arm, and 141 of these (60 %) retested at home. The majority of home arm retesters were comfortable having the kit posted to their home (86 %); found it easy to follow the instructions and collect the specimens (96 %); were confident they had collected the specimens correctly (90 %); and reported no problems (70 %). Most (65 %) preferred home retesting, 21 % had no preference and 14 % preferred clinic retesting. Comparing those with a preference for home testing to those who didn't, there were significant differences in being comfortable having a kit sent to their home ($p = 0.045$); not having been diagnosed with chlamydia previously ($p = 0.030$); and living with friends ($p = 0.034$). The overall cost for the home retest pathway was \$154 (AUD), compared to \$169 for the clinic-based retesting pathway and the cost per repeat infection detected was \$1409 vs \$3133.

Conclusions: Among individuals initially diagnosed with chlamydia in a sexual health clinic setting, home-based retesting was shown to be highly acceptable, preferred by most participants, and cost-efficient. However some clients preferred clinic-based testing, often due to confidentiality concerns in their home environment. Both options should be provided to maximise retesting rates.

Self-Collection

UriSwab™: an Effective Transport Medium for Nucleic Acid Detection of *Chlamydia Trachomatis*, *Mycoplasma Genitalium* And *Neisseria Gonorrhoeae*



Anna-Maria G. Costa^{1,2}, Suzanne M. Garland^{1,2,3,4}, Rebecca Guy⁵, Handan Wand⁵, Sepehr N. Tabrizi^{1,2,3,4}

Affiliation: ¹ Department of Microbiology and Infectious Diseases, The Royal Women's Hospital, Parkville, Australia

² Department of Microbiology, The Royal Children's Hospital, Parkville, Australia ³ Murdoch Childrens Research Institute, Parkville, Australia ⁴ Department of Obstetrics and Gynaecology, The University of Melbourne, Parkville, Australia

⁵ The Kirby Institute, University of New South Wales, Sydney, Australia

Keywords

FLOQSwabs®

Chlamydia trachomatis

Neisseria gonorrhoeae

Mycoplasma genitalium

Abstract

Background: Patient self-sampling allows for remote collection and return to clinic or laboratory by post. Urine samples, although convenient, are challenging to post. This study evaluated UriSwab™ (Copan, Brescia, Italy) as a collection and transport vessel for *Chlamydia trachomatis* (CT), *Neisseria gonorrhoeae* (NG) and *Mycoplasma genitalium* (MG) detection by polymerase chain reaction, compared with flocked swab and neat urine.

Methods: Five replicates of each specimen type were prepared from previously characterised urine samples (n = 330), stored at room temperature (RT) or 37°C, then extracted on day 1, 3, 7, 10 and 16 (VERSANT kPCR Sample Prep System, Siemens, Munich, Germany). Crossing thresholds (Cq) from CT and NG detection (VERSANT CT/GC DNA 1.0 assay kit, Siemens) and MG detection (real-time polymerase chain reaction assay) were compared using logistic regression, stratified by sample type, temperature and analyte. Mixed-model statistical techniques were used to assess correlation between repeated observations.

Results: UriSwab™ showed an increasing trend in Cq values at RT and 37°C for CT and NG, and RT for MG (all P < 0.01). UriSwab™ was not statistically significantly different to neat urine, except CT at RT (0.83, 95% confidence interval: 0.51–1.15). Flocked swab similarly showed increasing Cq values at 37°C for CT, a significant decreasing trend at RT for MG and increasing trend at 37°C for MG. Flocked swab was not statistically significantly different from neat urine at RT and 37°C for CT and MG. **Conclusion:** UriSwab allows transport of urine for CT, NG and MG detection regardless of storage time or temperature, suggesting that CT and NG are stable for up to 16 days and MG up to 10 days.

Self-Collection

A Randomized Comparison of Different Vaginal Self-sampling Devices and Urine for Human Papillomavirus Testing—Predictors 5.1



Louise Cadman¹, Caroline Reuter¹, Mark Jitlal¹, Michelle Kleeman^{1,2}, Janet Austin¹, Tony Hollingworth¹, Anna L. Parberry³, Lesley Ashdown-Barr¹, Deepali Patel^{1,2}, Belinda Nedjai¹, Attila T. Lorincz¹, Jack Cuzick¹
Affiliation: 1 Centre for Cancer Prevention, Wolfson Institute of Preventive Medicine, Queen Mary University of London, London, United Kingdom. 2 NIHR Biomedical Research Centre, Guy's and St Thomas' NHS Foundation Trust and King's College, London, United Kingdom. 3 Colposcopy Department, Royal London Hospital, London, United Kingdom.
Journal: Cancer Epidemiology, Biomarkers & Prevention (2021)

Keywords

Copan FLOQSwabs[®]

HPV

Vaginal sampling

Abstract

Background: Human papillomavirus (HPV)-based screening is rapidly replacing cytology as the cervical screening modality of choice. In addition to being more sensitive than cytology, it can be done on self-collected vaginal or urine samples. This study will compare the high-risk HPV positivity rates and sensitivity of self-collected vaginal samples using four different collection devices and a urine sample. Methods: A total of 620 women referred for colposcopy were invited to provide an initial stream urine sample collected with the Colli-Pee device and take two vaginal self-samples, using either a Copan dry flocced swab (DF) and a wet dacron swab (WD), or a HerSwab (HS) and Qvintip (QT) device. HPV testing was performed by the BD Onclarity HPV Assay. Results: A total of 600 vaginal sample pairs were suitable for analysis, and 505 were accompanied by a urine sample. Similar positivity rates and sensitivities for CIN2⁺ and CIN3⁺ were seen for DF, WD, and urine, but lower values were seen for QT and HS. No clear user preferences were seen between devices, but women found urine easiest to collect, and were more confident they had taken the sample correctly. The lowest confidence in collection was reported for HS. Conclusions: Urine, a DF swab, and WD swab all performed well and were well received by the women, whereas the Qvintip and HerSwab devices were less satisfactory. This is the first study to compare five self-sampling methods in the same women taken at the same time. It supports wider use of urine or vaginal self-sampling for cervical screening.

Self-Collection

The feasibility and acceptability of self-sampling and HPV testing using Cepheid Xpert® HPV in a busy primary care facility



Authors: YL Woo¹

Affiliation: 1 Faculty of Medicine, University of Malaysia on behalf of ROSE team

Keywords

Copan FLOQSwabs®

HPV

Vaginal sampling

ROSE Project

Abstract

Malaysia's approach to reducing the burden of HPV-related disease has centred on adolescent vaccination and cervical screening with Pap smears. While the vaccination programme has been broadly successful, Pap smear screening has been less successful. In an effort to improve screening uptake, the ROSE 1.0 pilot aimed to create more efficient screening, with improved quality and lower total cost. The outcome of this design process was a new cervical screening approach, based on self-sampling using Copan FLOQswabs™ (Copan, Brescia, Italy) in community clinics (Klinik Kesihatan) and the Cepheid Xpert® HPV assay system, with testing performed on site. Advantages of this switch from Pap smear to HPV DNA testing include improved sensitivity (90% compared with 50%) and an automated, objective process enabling greater cost effectiveness.

Self-Collection

Randomized trial evaluating self-sampling for HPV DNA based tests for cervical cancer screening in Nigeria



Authors: Fatima Modibbo¹, K. C. Iregbu¹, James Okuma², Annemiek Leeman³, Annemieke Kasius³, Maurits de Koning³, Wim Quint³, Clement Adebamowo^{4,5}

Affiliations: 1 Department of Medical Microbiology and Parasitology, National Hospital Abuja, Plot 132 Central Business District (Phase II), PMB 425 Garki, Abuja 90001, Nigeria. 2 Department of Research, Institute of Human Virology, Abuja, Nigeria. 3 DDL Diagnostic Laboratory, Visseringlaan 25, 2288, ER, Rijswijk, Netherlands. 4 Department of Epidemiology and Public Health, University of Maryland School of Medicine, Baltimore, MD, USA. 5 Institute of Human Virology and Greenebaum Comprehensive Cancer Centre, University of Maryland School of Medicine, Baltimore, MD, USA.

Keywords

Copan FLOQSwabs®

HPV

Cervical cancer screening

Randomized trial

Abstract

Background: Cervical cancer incidence and mortality rates in Sub-Saharan Africa (SSA) remain high due to several factors including low levels of uptake of cervical cancer screening. Self-collection of cervicovaginal samples for HPV DNA testing may be an effective modality that can increase uptake of cervical cancer screening in SSA and hard to reach populations in developed countries. We investigated whether self-collection of cervicovaginal samples for HPV DNA tests would be associated with increased uptake of screening compared with clinic-based collection of samples. Furthermore, we compared the quality of samples collected by both approaches for use in HPV genotyping. Methods: We conducted a community based randomized trial in a semi-urban district of Abuja, Nigeria with 400 women, aged 30 to 65 years randomized to either hospital-collection or self-collection of cervicovaginal swab samples (Copan Italia). We compared cervical cancer screening uptake among the 2 groups and evaluated the concentration of human DNA in the samples by measuring RNase P gene levels using qPCR. High-risk HPV DNA detection and typing was done using the GP5+/6+ Luminex system. Results: Most participants in the self-collection arm (93%, 185/200) submitted their samples while only 56% (113/200) of those invited to the hospital for sample collection attended and were screened during the study period (p value < 0.001). Human genomic DNA was detected in all but five (1.7%) participants, all of whom were in the self-collection arm. The prevalence of high-risk HPV in the study population was 10% with types 35, 52 and 18 being the commonest. Conclusions: Our study shows that self-sampling significantly increased uptake of HPV DNA based test for cervical cancer screening in this population and the samples collected were adequate for HPV detection and genotyping. Cervical cancer screening programs that incorporate self-sampling and HPV DNA tests are feasible and may significantly improve uptake of cervical cancer screening in SSA.

Self-Collection

Acceptability of human papillomavirus (HPV) self-sampling among never- and under-screened Indigenous and other minority women: a randomised three-arm community trial in Aotearoa New Zealand



Authors: Naomi Brewer¹, Karen Bartholomew², Jane Grant², Anna Maxwell², Georgina McPherson³, Helen Wihongi², Collette Bromhead⁴, Nina Scott⁵, Sue Crengle⁶, Sunia Foliaki¹, Chris Cunningham¹, Jeroen Douwes¹, John D. Potter^{1,7}
Affiliations: 1 Research Centre for Hauora and Health, Massey University, PO Box 756, Wellington 6140, New Zealand. 2 Waitemata District Health Board (DHB) and Auckland DHB, Private Bag 93-503, Takapuna, Auckland 0740, New Zealand. 3 Waitemata District Health Board, Auckland, New Zealand. 4 School of Health Sciences, Massey University, Wellington, New Zealand. 5 University of Auckland, Waikato District Health Board, New Zealand. 6 Department of Preventive and Social Medicine, University of Otago, Dunedin, New Zealand. 7 Fred Hutchinson Cancer Research Center, Seattle, WA, USA

Keywords

Copan FLOQSwabs®

HPV

Minorities

HPV DNA testing

Abstract

Background: Internationally, self-sampling for human papillomavirus (HPV) has been shown to increase participation in cervical-cancer screening. In Aotearoa New Zealand, there are long-standing ethnic inequalities in cervical-cancer screening, incidence, and mortality, particularly for indigenous Maori women, as well as Pacific and Asian women.
Methods: We invited never- and markedly under-screened (≥ 5 years overdue) 30-69-year-old Maori, Pacific, and Asian women to participate in an open-label, three-arm, community-based, randomised controlled trial, with a nested sub-study. We aimed to assess whether two specific invitation methods for swab self-sampling (Copan Italia) improved screening participation over usual care among the least medically served populations. Women were individually randomised 3:3:1 to: clinic-based self-sampling (CLINIC – invited to take a self-sample at their usual general practice); home-based self-sampling (HOME – mailed a kit and invited to take a self-sample at home); and usual care (USUAL – invited to attend a clinic for collection of a standard cytology sample). Neither participants nor research staff could be blinded to the intervention. In a subset of general practices, women who did not participate within three months of invitation were opportunistically invited to take a self-sample, either next time they attended a clinic or by mail.
Findings: We randomised 3,553 women: 1,574 to CLINIC, 1,467 to HOME, and 512 to USUAL. Participation was highest in HOME (14.6% among Maori, 8.8% among Pacific, and 18.5% among Asian) with CLINIC (7.0%, 5.3% and 6.9%, respectively) and USUAL (2.0%, 1.7% and 4.5%, respectively) being lower. In fully adjusted models, participation was statistically significantly more likely in HOME than USUAL: Maori OR=9.7, (95%CI 3.0-31.5); Pacific OR=6.0 (1.8-19.5); and Asian OR=5.1 (2.4-10.9). There were no adverse outcomes reported. After three months, 2,780 non-responding women were invited to participate in a non-randomised, opportunistic, follow-on substudy. After 6 months, 192 (6.9%) additional women had taken a self-sample.

Self-Collection

The Acceptability and Preference of Vaginal Self-sampling for Human Papillomavirus (HPV) Testing among a Multi-ethnic Asian Female Population



Authors: Su Pei Khoo¹, Wen Tzian Lim¹, Reena Rajasuriar², Nazrila Hairizan Nasir³, Patti Gravitt⁴, Yin Ling Woo¹
Affiliations: 1 Department of Obstetrics and Gynaecology, Faculty of Medicine, University of Malaya, Kuala Lumpur Malaysia. 2 Department of Medicine, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia. 3 Division of Family Health Development, Ministry of Health, Putrajaya, Malaysia. 4 Department of Epidemiology and Public Health, University of Maryland School of Medicine, Baltimore, Maryland.

Keywords

Copan FLOQSwabs®

HPV

Acceptability

Self-sampling

Abstract

Abstract: Vaginal self-sampling for human papillomavirus (HPV) testing can potentially increase cervical screening coverage. This study aimed to investigate the acceptability of vaginal self-sampling for HPV testing and factors that might influence a woman's preference for this as a cervical screening method. This was a cross-sectional study that recruited 725 women from the urban and suburban areas of Selangor, Malaysia. All study participants were instructed to self-collect vaginal sample using a dry flocced swab (Copan Italia) before responding to a detailed questionnaire documenting their experience and preference for self-sampling. Most of the study participants (>80%) perceived vaginal self-sampling as easy, convenient, not embarrassing, comfortable, and were confident in performing the test. This suggests high acceptability toward vaginal self-sampling for HPV testing. Of the 725 women, 83% preferred self-sampling HPV testing over healthcare personnel sampling HPV testing and Pap test. Women with higher household income and full-time employment status were more likely to prefer self-sampling. Those who had not undergone Pap test also expressed preference for self-sampling HPV testing. Convenience and women's confidence in performing a vaginal self-sampling for HPV testing were the independent key factors that influenced the preference for self-sampling method. Vaginal self-sampling for HPV testing is highly acceptable among Malaysian women. It is the preferred choice as a primary cervical screening method and serves as an alternative to healthcare-acquired sample for Pap test.

Self-Collection

Implementing the 3T-approach for cervical cancer screening in Cameroon: Preliminary results on program performance



Authors: Juliette Levy¹, Marie de Preux¹, Bruno Kenfack², Jessica Sormani^{3,4}, Rosa Catarino³, Eveline F. Tinchos, Chloé Frund³, Jovanny T. Fouogue⁶, Pierre Vassilakos^{3,7}, Patrick Petignat³

Affiliations: 1 Faculty of Medicine, University of Geneva, Geneva, Switzerland. 2 Faculty of Medicine and Pharmaceutical Sciences, University of Dschang, Dschang District Hospital, Dschang, Cameroon. 3 Gynecology Division, Department of Gynecology and Obstetrics, University Hospitals of Geneva, Geneva, Switzerland. 4 Geneva School of Health Sciences, HESSO University of Applied Sciences and Arts Western Switzerland, Geneva, Switzerland. 5 Faculty of Medicine and Biomedical Sciences, Centre Hospitalier Universitaire (CHUY), Yaoundé, Cameroon. 6 Department of Obstetrics and Gynecology, Bafoussam Regional hospital, Bafoussam, Cameroon. 7 Geneva Foundation for Medical Education and Research, Genève, Switzerland

Keywords

Copan FLOQSwabs®

HPV

Screen-and-treat

Management

Abstract

Option recommended by World Health Organization (WHO) includes human papillomavirus (HPV) primary screening followed by visual inspection with acetic acid (VIA) triage. We implemented a program based on a 3T-approach (Test-Triage and Treat). Our objective was to verify the effectiveness of the program by defining a set of performance indices. A sensitization campaign was performed in Dschang (Cameroon) and women aged 30-49 years were invited to participate for screening based on the 3T-approach. Participants performed HPV swab self-sampling (Copan Italia), analysed with the point-of-care Xpert HPV assay followed by VIA/VILI triage and treatment if required. Key performance indicators (KPIs) for screening, diagnosis, treatment and follow-up were defined, and achievable targets were described for which the approach is likely to be running optimally. A total of 840 women with a mean age of 39.4±5.9 years participated. The KPIs included (i) the screening rate (8.4% at 7 months, target =20% at 12 months), (ii) HPV positivity rate (19.8%, expected range 18-25%), (iii) compliance to referral to VIA/VILI and complete test (100%, target >90%), (iv) compliance to referral to thermal ablation (100%, target > 90%), (v) VIA/VILI positivity rate (50.6%, expected range 45-55%), (vi) a single visit from diagnostic to treatment (79.8%, target >80%), (vii) compliance to follow up at 1 month (96.4%, target >80%) and (viii) at 6 months (70.6%, target >80%). Program performance based on the single-visit 3T-approach corresponded to defined targets and preliminary results support adequateness of KPIs for periodic monitoring.

Self-Collection

Exploring Factors Associated with Patients Who Prefer Clinician-Sampling to HPV Self-Sampling: A Study Conducted in a Low-Resource Setting



Authors: Jessica Sormani^{1,2}, Bruno Kenfack^{3,4}, AniaWisniak¹, Alida Moukam Datchoua⁴, Sophie Lemoupa Makajio^{1,5}, Nicole C. Schmidt^{1,6}, Pierre Vassilakos^{1,7}, Patrick Petignat¹

Affiliations:1 Gynaecology Division, Department of Paediatrics, Gynaecology and Obstetrics, University Hospitals of Geneva, 1205 Geneva, Switzerland. 2 School of Health Sciences, HES-SO University of Applied Sciences and Arts Western Switzerland, 1227 Geneva, Switzerland. 3 Department of Obstetrics and Gynecology, Faculty of Medicine and Pharmaceutical Sciences, University of Dschang, Dschang, Cameroon. 4 Department of Gynaecology and Obstetrics, District Hospital of Dschang, Dschang, Cameroon. 5 Faculty of Medicine, Institute of Global Health, University of Geneva, 1205 Geneva, Switzerland. 6 Faculty of Social Science, Catholic University of Applied Science, 55122 Mainz, Germany. 7 Geneva Foundation for Medical Education and Research, 1202 Geneva, Switzerland

Keywords

Copan FLOQSwabs®

HPV

Low-Resource Setting

Self-sampling

Abstract

Human papillomavirus (HPV) self-sampling (Self-HPV) is a promising strategy to improve cervical cancer screening coverage in low-income countries. However, issues associated with women who prefer conventional HPV clinical-sampling over HPV self-sampling may affect screening participation. To address this issue, our study assessed factors associated with women's preferences related to Self-HPV. This study was embedded in a large clinical trial recruiting women aged 30–49 years in a primary HPV-based study termed "3T-Approach" (for Test-Triage-Treatment), launched in 2018 at Dschang District Hospital, West Cameroon. Participants were invited to perform a Self-HPV. After the sampling and before receiving the results, participants completed a questionnaire about cervical cancer screening and their preferences and perceptions around Self-HPV. The median age of the 2201 participants was 40.6 (IQR 35–45) years. Most (1693 (76.9%)) preferred HPV self-sampling or had no preference for either method, and 508 (23.1%) preferred clinician-sampling. Factors associated with an increased likelihood of reporting a clinician-sampling preference were tertiary educational level (29.4% CI: 25.6–33.6 vs. 14.4% CI: 12.8–16.1) and being an employee with higher grade professional or managerial occupations (5.5% CI: 3.8–7.9 vs. 2.7% CI: 2.0–3.5). The main reported reason for women preferring clinician-sampling was a lack of "self-expertise". Most women (>99%) would agree to repeat HPV self-sampling and would recommend it to their relatives. HPV self-sampling in the cultural context of central Africa was well accepted by participants, but some participants would prefer to undergo clinician sampling. Health systems should support well-educated women to increase self-confidence in using HPV self-sampling.

Self-Collection

Non-speculum sampling approaches for cervical screening in older women: randomised controlled trial



Authors: Rebecca Landy¹, Tony Hollingworth², Jo Waller³, Laura AV Marlow³, Jane Rigney³, Thomas Round⁴, Peter D Sasieni³, Anita WW Lim³

Affiliations: 1 Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Department of Health and Human Services, Bethesda, Maryland, US. 2 Whipps Cross University Hospital, Barts Health NHS Trust, London. 3 Comprehensive Cancer Centre, School of Cancer and Pharmaceutical Sciences, Faculty of Life Sciences and Medicine, King's College London, London. 4 School of Population Health and Environmental Sciences, King's College London, London and National Cancer Analysis and Registration Service, Public Health England.

Keywords

Copan FLOQSwabs®

Human papillomavirus
DNA test

older women

Self sampling

Abstract

Background: Cervical cancer disproportionately affects women ≥ 65 years, especially those not screened regularly. Speculum use is a key barrier. Aim: To assess if offering non-speculum clinician taken sampling and self-sampling increases uptake for lapsed attenders aged 50–64 years. Design and setting: Pragmatic randomised control trial conducted at 10 general practices in East London, UK. Method: Participants were 784 women aged 50–64 years, last screened 6–15 years before randomisation. Intervention participants received a letter offering the choice of non-speculum clinician- or self-sampling. Control participants received usual care. The main outcome measure was uptake within 4 months. Results: Screening uptake 4 months after randomisation was significantly higher in the intervention arm: 20.4% (n = 80/393) versus 4.9% in the control arm (n = 19/391, absolute difference 15.5%, 95% confidence interval [CI] = 11.0% to 20.0%, $P < 0.001$). This was maintained at 12 months: intervention 30.5% (n = 120/393) versus control 13.6% (n = 53/391) (absolute difference 17.0%, 95% CI = 11.3% to 22.7%, $P < 0.001$). Conventional screening attendance within 12 months was very similar for both intervention 12.7% (n = 50/393) and control 13.6% (n = 53/391) arms. Ethnic differences were seen in screening modality preference. More White women opted for self-sampling (50.7%, n = 38/75), whereas most Asian and Black women and those from other ethnic backgrounds opted for conventional screening. Conclusion: Offering non-speculum clinician-taken sampling and self-sampling substantially increases uptake in older lapsed attendee women. Nonspeculum clinician sampling appeals to women who dislike the speculum but still prefer a clinician to take their sample. Providing a choice of screening modality may be important for optimising cervical screening uptake.

Self-Collection

Comparison of Different Self Sampling Devices for Molecular Detection of Human Papillomavirus (HPV) and Other Sexually Transmitted Infections (STIs): A Pilot Study



Authors: Illari Sechi¹, Cocuzza Clementina Elvezia², Marianna Martinelli², Narcisa Muresu¹, Santina Castriciano³, Giovanni Sotgiu⁴, Andrea Piana¹

Affiliations: 1 Department of Medical, Surgical and Experimental Sciences, University of Sassari, Sassari, Italy. 2 Department of Medicine and Surgery, University of Milano-Bicocca, Monza, Italy. 3 Copan Italia Spa, Brescia, Italy. 4 Clinical Epidemiology and Medical Statistics Unit, Department of Medical, Surgical and Experimental Sciences, University of Sassari, Sassari, Italy

Keywords

Copan FLOQSwabs®

HPV

sexually transmitted infections

Self-collected samples

Abstract

Background: Cervical cancer is the fourth most common cancer in women, and it is well known that high-risk human papillomavirus (hrHPV) infections are the necessary carcinogenic factors for the development of cervical tumors. Moreover, the interaction between HPV and other sexually transmitted infections (STIs) may increase the risk of cancer progression. Self-sampling has been demonstrated to represent a valid and well-accepted alternative, favoring women's participation in screening programs. This study aimed to investigate the use of FLOQSwabs® (FS) as compared to two other vaginal self-collection devices for the detection of hrHPV and other sexually transmitted infections. **Methods:** Cervical and vaginal self-samples were collected, using two different combinations of vaginal self-sampling devices, from 40 women referred to colposcopy for a documented abnormal Pap smear. All samples were tested for hrHPV and seven STI pathogens using two commercial molecular assays. **Results:** Data on hrHPV detection from the first group of women showed an almost perfect agreement (kappa: 0.89) between cervical vs. FS vaginal self-samples, and a substantial agreement (kappa: 0.79) between cervical and HerSwab™ (HS) samples. In the second group of women, an almost perfect agreement (kappa: 0.90) was demonstrated in the detection of hrHPV between cervical samples vs. FS, and a moderate agreement (kappa: 0.60) for cervical vs. Evalyn®Brush (EB) self-collected samples. STI detections showed a very good agreement (kappa: 0.89 and kappa: 1.00) both among FS vs. HS and FS vs. EB, respectively. There was no statistically significant difference between the different devices used. The most frequently detected hrHPV genotypes in the studied population were HPV 16, 31, 35, 51, and 56; whilst the most frequently identified STI pathogens were *Ureaplasma parvum* and *Mycoplasma hominis*. Overall, investigated women did not report any discomfort in using the different vaginal self-collection devices. **Conclusion:** Evaluation of the three different vaginal self-collection devices confirmed their overall good acceptability by the studied population, as well as a similar agreement for hrHPV detection as compared to cervical samples. Our study indicated that the use of self-collected samples offers an alternative strategy to improve women's participation in cervical cancer screening programs, but also underlined the importance of evaluating the concordance in hrHPV detection of collection devices in combination with the molecular hrHPV assay.

Self-Collection

Saliva detection of SARS-CoV-2 for mitigating company outbreaks: a surveillance experience, Milan, Italy, March 2021



Authors: Emerenziana Ottaviano¹, Chiara Parodi¹, Elisa Borghi¹, Valentina Massa¹, Cristina Gervasini¹, Stefano Centanni¹, Gianvincenzo Zuccotti², LollipopStudy Group, Silvia Ancona, Elisa Adele Colombo, Elisabetta Di Fede, Paolo Graziosi, Elena Lesma, Antonella Lettieri, Silvia Bianchi¹

Affiliations: 1 Department of Health Sciences, Università degli Studi di Milano, Milan, Italy. 2 Department of Paediatrics, Children Hospital V. Buzzi, Università degli Studi di Milano, Milan, Italy

Keywords

Copan Lollisponge™

Saliva sample

SARS-CoV-2

Self-collection

Abstract

Monitoring the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) communitywide transmission with a suitable and effective sampling method would be of great support for public health response to the spreading due to asymptomatic subjects in the community. Here, we describe how using saliva samples, collected with Copan Lollisponge™, for SARS-CoV-2 detection has allowed for a weekly surveillance of a small business company and the early detection of coronavirus disease 2019 cases. As on 23rd March, two cases were detected and investigated, and control measures were rapidly applied.



This document may contain product information otherwise not accessible or valid in your country. Please be aware that Copan Italia S.p.A. does take any responsibility for accessing such information which may not comply with any valid legal process, regulation, registration or usage in the country of your origin. Product clearance and availability restrictions may apply in some Countries. Please refer to Copan website (www.copangroup.com) to view and/or download the most recent version of the brochure. This document is mainly intended for marketing purposes, always consult product insert for complete information. The original poster/study has been amended and/or some portions omitted for editorial reasons. ©2021 Copan Italia. All rights reserved. The trademarks mentioned herein are property of Copan Italia S.p.A.
Code: JMKR012R01



@copangroup

Copan Italia s.p.a.
Via Francesco Perotti 10,
25125 Brescia, Italy

t | f +030 2687211
@ | info@copangroup.com
www.copangroup.com